

Dense and narrow rings discovered around the Centaur object (10199) Chariklo

B. Sicardy¹, F. Braga-Ribas², G. Benedetti-Rossi¹, N. Ligier¹, L. Maquet¹, F. Roques¹, R. Leiva Espinoza³, M. El Moutamid⁴, R. Duffard⁵, N. Morales⁵, J.L. Ortiz⁵, C. Snodgrass⁶, J. Skottfelt⁷, K. Harpsøe⁷, U. G. Jørgensen⁷, E. Jehin⁸, J. Pollock⁹, M. Assafin¹⁰, J. Camargo², J. Desmars², R. Vieira-Martins², C. Dumas¹¹, V. Ivanov¹¹, A. Maury¹², H. Korhonen¹³, and M. Leitzinger¹⁴

¹LESIA, Observatoire de Paris, CNRS UMR 8109, Université Pierre et Marie Curie, Université Paris-Diderot, Meudon, France

²Observatorio rio Nacional/MCTI Rio de Janeiro, Brazil

³Instituto de Astrofísica, Facultad de Física, Pontificia Universidad Católica de Chile, Santiago, Chile

⁴Department of Astronomy, Cornell University, Ithaca, USA

⁵Instituto de Astrofísica de Andalucía, Granada, Spain

⁶Max Planck Institute for Solar System Research, Goettingen, Germany

⁷Niels Bohr Institute & Geological Museum, Copenhagen, Denmark

⁸Institut d'Astrophysique, Université de Liège, Belgium

⁹Physics and Astronomy Department, Appalachian State University, Boone, USA

¹⁰Observatorio do Valongo/UFRJ, Rio de Janeiro, RJ, Brazil

¹¹European Southern Observatory, Santiago, Chile

¹²San Pedro de Atacama Celestial Explorations, San Pedro de Atacama, Chile

¹³Finnish Centre for Astronomy with ESO, University of Turku, Piikkiö, Finland

¹⁴University of Graz, Institute of Physics, Department for Geophysics, Astrophysics, and Meteorology/IGAM, Graz, Austria

A stellar occultation observed on June 3, 2013 revealed the unexpected presence of two dense rings around (10199) Chariklo [1], the largest Centaur object known to date with a radius of 124 ± 9 km [2]. The two rings have respective orbital radii, widths, and normal optical depths of $a_1 = 391$ km, $W_1 = 7$ km, $\tau_1 = 0.4$ and $a_2 = 405$ km, $W_2 = 3$ km, $\tau_2 = 0.06$ [1]. They are separated by a gap of about 9 km with an optical depth less than 0.004 (1-sigma limit). The presence of those rings has been confirmed during another stellar occultation observed from ESO/NTT La Silla, ESO/VLT Paranal and San Pedro de Atacama on February 16, 2014. More results on the azimuthal variations of the rings and mass estimations of their putative shepherding satellites will be presented.

This is the first ring system ever observed that does not pertain to a giant planet. The existence of such a system raises several questions as to the origin and evolution of rings around such a small object. This discovery also suggests that rings may be a more frequent feature than previously thought, in particular, around small bodies. Possible models for the ring formation will be proposed. They can be classified into collisional scenarios that disrupted an impactor or a pre-existing satellite, tidal disruption of an inward-migrating satellite, or material produced by a cometary activity of the central body.

Acknowledgements: The main author thanks support from the French grant 'Beyond Neptune II' Blanc Inter II SIMI 5–6.

References: [1] Braga-Ribas et al. (2014), Nature 26 March 2014, doi:10.1038/nature13155; [2] Fornasier, S. et al. (2013), Astron. Astrophys. 555, A15.